



Address 1: 2445 Country Place Blvd., Suite 103,  
New Port Richey, FL 34655

Address 2: 6928 W. Linebaugh Ave. Suit 102, Tampa,  
FL 33625

Phone: (813) 749-7143

Fax: (813) 264-9262

Website: [www.floridamedspas.com](http://www.floridamedspas.com)

## Post-Insertion Instructions for Women

- ‡ Your insertion site has been covered with two layers of bandages. Remove the outer pressure bandage any time after 48 hours. It must be removed as soon as it gets wet. You may replace it with a bandage to catch any anesthetic that may ooze out. The inner layer is either waterproof foam tape or steri-strips. They may be removed in **3 days**.
- ‡ We recommend putting an ice pack on the insertion area a couple of times for about 20 minutes each time over the next 4 to 5 hours.
- ‡ Do not take tub baths or get into a hot tub or swimming pool for **3 days**. You may shower but do not scrub the site until the incision is well healed (about 7 days).
- ‡ No major exercises for the incision area for the next **5 days**, this includes running, riding a horse, etc.
- ‡ The sodium bicarbonate in the anesthetic may cause the site to swell for 1-3 days.
- ‡ The insertion site may be uncomfortable for up to 2 to 3 weeks. If there is itching or redness you may take Benadryl for relief, 50 mg. orally every 6 hours. Caution this can cause drowsiness!
- ‡ You may experience bruising, swelling, and/or redness of the insertion site which may last from a few days up to 2 to 3 weeks.
- ‡ You may notice some pinkish or bloody discoloration of the outer bandage. This is normal.
- ‡ If you experience bleeding from the incision, apply firm pressure for 5 minutes.
- ‡ Please call if you have any bleeding not relieved with pressure (not oozing), as this is NOT normal — (813)749-7143
- ‡ Please call if you have any pus coming out of the insertion site, as this is NOT normal — (813)749-7143

### Reminders:

- ‡ Remember to go for your post-insertion blood work 5-6 weeks after the insertion. ‡ Most women will need re-insertions of their pellets 4-5 months after their initial insertion.
- ‡ Please call as soon as symptoms that were relieved from the pellets start to return to make an appointment for a re-insertion. The charge for the second visit will only be for the insertion and not a consultation.

### Additional Instructions:

I acknowledge that I have received a copy and understand the instructions on this form.

Print Name

Signature

Today's Date



Address 1: 2445 Country Place Blvd., Suite 103,  
New Port Richey, FL 34655

Address 2: 6928 W. Linebaugh Ave. Suit 102, Tampa,  
FL 33625

Phone: (813) 749-7143

Fax: (813) 264-9262

Website: [www.floridamedspas.com](http://www.floridamedspas.com)

## What Might Occur After A Pellet Insertion

A significant hormonal transition will occur in the first four weeks after the insertion of your hormone pellets. Therefore, certain changes might develop that can be bothersome.

- ‡ **FLUID RETENTION:** Testosterone stimulates the muscle to grow and retain water, which may result in a weight change of two to five pounds. This is only temporary. This happens frequently with the first insertion, and especially during hot, humid weather conditions.
- ‡ **SWELLING OF THE HANDS & FEET:** This is common in hot and humid weather. It may be treated by drinking lots of water, reducing your salt intake, taking cider vinegar capsules daily, (found at most health and food stores) or by taking a mild diuretic, which the office can prescribe.
- ‡ **UTERINE SPOTTING/BLEEDING:** This may occur in the first few months after an insertion, especially if you have been prescribed progesterone and are not taking properly: i.e. missing doses, or not taking a high enough dose. Please notify the office if this occurs. Bleeding is not necessarily an indication of a significant uterine problem. More than likely, the uterus may be releasing tissue that needs to be eliminated. This tissue may have already been present in your uterus prior to getting pellets and is being released in response to the increase in hormones.
- ‡ **MOOD SWINGS/IRRITABILITY:** These may occur if you were quite deficient in hormones. They will disappear when enough hormones are in your system. 5HTP can be helpful for this temporary symptom and can be purchased at many health food stores.
- ‡ **FACIAL BREAKOUT:** Some pimples may arise if the body is very deficient in testosterone. This lasts a short period of time and can be handled with a good face cleansing routine, astringents and toner. If these solutions do not help, please call the office for suggestions and possibly prescriptions.
- ‡ **HAIR LOSS:** Is rare and usually occurs in patients who convert testosterone to DHT. Dosage adjustment generally reduces or eliminates the problem. Prescription medications may be necessary in rare cases.
- ‡ **HAIR GROWTH:** Testosterone may stimulate some growth of hair on your chin, chest, nipples and/or lower abdomen. This tends to be hereditary. You may also have to shave your legs and arms more often. Dosage adjustment generally reduces or eliminates the problem.

I acknowledge that I have received a copy and understand the instructions on this form.

Print Name

Signature

Today's Date



Address 1: 2445 Country Place Blvd., Suite 103,  
New Port Richey, FL 34655  
Address 2: 6928 W. Linebaugh Ave. Suit 102, Tampa,  
FL 33625  
Phone: (813) 749-7143  
Fax: (813) 264-9262  
Website: www.floridamedspas.com

## Female Treatment Plan

- 1 The following medications or supplements are recommended in addition to your pellet therapy.
- 2 Please refer to the supplement brochure to help you understand why these are beneficial.
- 3 Unless specified, these can be taken anytime of day without regards to meals.

**Supplements:** These may be purchased in our office. When you run out they can be mailed to you for your convenience.

\_\_\_\_\_ ADK (vitamins A, D3 and K2)  
\_\_\_\_\_ 1 a day                      \_\_\_\_\_ 2 a day for weeks, then one a day  
\_\_\_\_\_ BioTE Iodine Plus 12.5 mg daily with food

2000-4000mg Vitamin C, B-complex and 1/2 tsp Celtic or Mediterranean sea salt mixed in warm water daily are recommended for the first few weeks you are taking BioTE Iodine Plus to prevent potential detox symptoms (headache, body aches, fatigue, irritability)

\_\_\_\_\_ DIM 150 mg daily- increases free testosterone levels while naturally decreasing "bad" estrogen

**Prescriptions: These have been called into your preferred pharmacy**

\_\_\_\_\_ Progesterone/Prometrium nightly  
\_\_\_\_\_ 100 mg \_\_\_\_\_ 200 mg

Please do not skip doses of this medication as it can result in vaginal bleeding or an increased risk for endometrial cancer.

\_\_\_\_\_ Nature Throid \_\_\_\_\_ mg every morning. This should be taken on an empty stomach. Please wait 30 minutes before putting anything else on your stomach. This includes coffee, food, medications, vitamins or supplements.

\_\_\_\_\_ Sample given

\_\_\_\_\_ Wean of Synthroid/levothyroxine: alternate your desiccated thyroid (natur-throid) every other day with Synthroid/levothyroxine for 3 weeks then go to every day on your desiccated thyroid.

\_\_\_\_\_ Spironolactone 100 mg daily \_\_\_\_\_ (other)

\_\_\_\_\_ Wean off your antidepressant (see wean protocol) \_\_\_\_\_ (other)

**Please call or email for any questions about these recommendations.**

I acknowledge that I have received a copy and understand the instructions on this form.

Print Name

Signature

\_\_\_\_\_  
Today's Date



6928 W. Linebaugh Ave. Suit 102, Tampa, FL 33625  
 (813) 749-7143  
 (813) 264-9262  
 www.westchasespa.com

## OFFICE USE ONLY - Female Intake Form

Name:		Date:	
Height:	Weight:		
Current Medications:			
<b>Surgery/ History:</b> Hysterectomy: <input type="checkbox"/> Yes <input type="checkbox"/> No Ovaries: <input type="checkbox"/> Yes <input type="checkbox"/> No			
Last Pap:		Last Mammogram: Normal? <input type="checkbox"/> Yes <input type="checkbox"/> No	
<b>Symptoms:</b>			
<b>Labs:</b>			
Estradiol:	Testosterone:	FSH:	Vitamin D:
Vitamin B12:	TSH:	Free T3:	TPO: CBC:
Chem Panel:	LDL:	HDL:	Triglycerides:
<b>Plan:</b>			
<p>This patient presents today for hormone pellets. The procedure, risks, benefits and alternatives were explained to the patient. Questions were answered and a consent form for the insertion of Testosterone and/or Estradiol pellet implants was signed. An area in the hip was prepped with Betadine swabs. A sterile drape was applied. 1% Lidocaine with epinephrine and sodium bicarbonate was injected to anesthetize the area. A small transverse incision was made using a number 11 blade. The trocar with cannula was passed through the incision into the subcutaneous tissue. Testosterone and or Estradiol pellet(s) were inserted through the cannula into the subcutaneous tissue. Bleeding was minimal. Steri-strips and/or Foam Tape were applied. A sterile dressing was applied. The patient tolerated the procedure well. Postoperative instructions were reviewed and a copy given to the patient. Pellets used are as follows:</p>			
<b>Treat with:</b>			
<b>1. Testosterone:</b>		MG's	<b>Testosterone Lot Numbers:</b>
<b>2. Estradiol:</b>		MG's	<b>Estradiol Lot Numbers:</b>
Insertion Site: <input type="checkbox"/> Left Hip <input type="checkbox"/> Right Hip			
<b>3. Progesterone:</b>		Cycle or	<b>Continuous</b> (check one)
<b>4. Femara:</b>	<b>Arimidex:</b>	<b>DIM:</b>	
<b>5. Vitamin ADK:</b>	<b>Thyroid:</b>	<b>Iodine:</b>	
<b>6. Evening Primrose:</b>			
<b>7. Other:</b>			
<b>Comments:</b>			



6928 W. Linebaugh Ave. Suit 102, Tampa, FL 33625  
 (813) 749-7143  
 (813) 264-9262  
 www.westchasespa.com

## OFFICE USE ONLY - Female Patient Treatment Form

Name:	Date:
<b>Symptoms/Notes:</b>	
<b>Procedure Report:</b>	
<p>The procedure, risks, benefits and alternatives were explained to the patient. Questions were answered and a consent form for the insertion of Testosterone and/or Estradiol pellet implants was signed. An area in the hip was prepped with Betadine swabs. A sterile drape was applied. 1% Lidocaine with epinephrine and sodium bicarbonate was injected to anesthetize the area. A small transverse incision was made using a number 11 blade. The trocar with cannula was passed through the incision into the subcutaneous tissue. Testosterone and/or Estradiol pellet(s) were inserted through the cannula into the subcutaneous tissue. Bleeding was minimal. Steri-strips and/or Foam Tape were applied. A sterile dressing was applied. The patient tolerated the procedure well. Postoperative instructions were reviewed and a copy given to the patient.</p>	
Weight:	Estrogen pellet Lot #:
#:	Testosterone pellet Lot
Estradiol:	mg      Testosterone:      mg
Insertion site:	Left Hip      Right Hip
Date:	
<b>Symptoms/Notes:</b>	
<b>Procedure Report:</b>	



6928 W. Linebaugh Ave. Suit 102, Tampa, FL 33625  
 (813) 749-7143  
 (813) 264-9262  
 www.westchasespa.com

The procedure, risks, benefits and alternatives were explained to the patient. Questions were answered and a consent form for the insertion of Testosterone and/or Estradiol pellet implants was signed. An area in the hip was prepped with Betadine swabs. A sterile drape was applied. 1% Lidocaine with epinephrine and sodium bicarbonate was injected to anesthetize the area. A small transverse incision was made using a number 11 blade. The trocar with cannula was passed through the incision into the subcutaneous tissue. Testosterone and/or Estradiol pellet(s) were inserted through the cannula into the subcutaneous tissue. Bleeding was minimal. Steri-strips and/or Foam Tape were applied. A sterile dressing was applied. The patient tolerated the procedure well. Postoperative instructions were reviewed and a copy given to the patient.

Weight:	Estrogen pellet Lot #:		Testosterone pellet Lot
#:			
Estradiol:	mg	Testosterone:	mg
Insertion site:	Left Hip	Right Hip	